

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 1 022 332 A1

(12)

EUROPEAN PATENT APPLICATION

published in accordance with Art. 158(3) EPC

(43) Date of publication:
26.07.2000 Bulletin 2000/30

(51) Int. Cl.⁷: C12N 9/10, C12N 15/54,
C12N 1/19, C12P 19/00

(21) Application number: 98941819.9

(86) International application number:
PCT/JP98/04087

(22) Date of filing: 10.09.1998

(87) International publication number:
WO 99/13059 (18.03.1999 Gazette 1999/11)

(84) Designated Contracting States:
AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE

(30) Priority: 10.09.1997 JP 24515497

(71) Applicant:
Meiji Seika Kaisha, Ltd.
Tokyo 104-8002 (JP)

• NAKANE, Akitaka,
Meiji Seika Kaisha, Ltd.
Sakado-shi, Saitama 350-0289 (JP)
• KONO, Toshiaki,
Meiji Seika Kaisha, Ltd.
Sakado-shi, Saitama 350-0289 (JP)

(72) Inventors:
• YANAI, Koji,
Meiji Seika Kaisha, Ltd.
Sakado-shi, Saitama 350-0289 (JP)

(74) Representative:
Lewin, John Harvey et al
Elkington and Fife,
Prospect House,
8 Pembroke Road
Sevenoaks, Kent TN13 1XR (GB)

(54) BETA-FRUCTOFURANOSIDASE AND GENE THEREOF

(57) A novel β -fructofuranosidase and its gene are disclosed. A polypeptide comprising the amino acid sequence of SEQ ID No. 1 or No. 3 is an enzyme having β -fructofuranosidase activity and high transferase activity, and is capable of efficiently producing fructooligosaccharides.




FIG. 1A




FIG. 1B




FIG. 1C

EP 1 022 332 A1




FIG. 1D

EP 1 022 332 A1

Description

Background of the Invention5 Field of the Invention

[0001] The present invention relates to a β -fructofuranosidase having a fructose transferase activity, which is useful for the industrial production of fructooligosaccharides, and its gene and use.

10 Background Art

[0002] The molecular structure of a fructooligosaccharide is the same as that of sucrose, except that the fructose half of a fructooligosaccharide is coupled with another one to three fructose molecules at positions C1 and C2 via a β -bond. Fructooligosaccharides are indigestible sugars known for their physiological advantages, such as the facilitation of Bifidobacterial growth in the intestines, metabolic stimulation for cholesterol and other lipids, and little cariogenicity.

[0003] Fructooligosaccharides are found in plants, such as asparagus, onion, Jerusalem-artichoke and honey. They are also synthesized from sucrose by the newly industrialized mass production technique using fructosyltransfer reaction which is catalyzed by a β -fructofuranosidase derived from a microorganism.

[0004] The molecular structure of 1-kestose and nystose, which make up component of industrially produced fructooligosaccharide mixtures of today, are the same as that of sucrose except that their fructose half is coupled with one and two molecules of fructose, respectively. It has been found recently that their high-purity crystals exhibit new desirable characteristics both in physical properties and food processing purpose while maintaining the general physiological advantages of fructooligosaccharides (Japanese Patent Application No. 222923/1995, Japanese Laid-Open Publication No. 31160/1994). In this sense, they are fructooligosaccharide preparations having new features.

[0005] In consideration of the above, some of the inventors have already proposed an industrial process for producing crystal 1-kestose from sucrose (Japanese Patent Application No. 64682/1996, Japanese Patent Application No. 77534/1996, and Japanese Patent Application No. 77539/1996). According to this process, a β -fructofuranosidase harboring fructosyltransferase activity is first allowed to act on sucrose to produce 1-kestose; the resultant 1-kestose is fractionated to a purity of 80% or higher by chromatographic separation; then, using this fraction as a crystallizing sample, crystal 1-kestose is obtained at a purity of 95% or higher. The β -fructofuranosidase harboring fructosyltransferase activity used in this process should be able to produce 1-kestose from sucrose at a high yield while minimizing the byproduct nystose, which inhibits the reactions in the above steps of chromatographic separation and crystallization. In the enzyme derived from Aspergillus niger, which is currently used for the industrial production of fructooligosaccharides mixtures, the 1-kestose yield from sucrose is approximately 44%, while 7% is turned to nystose (Japanese Patent Application No. 64682/1996). These figures suggest that the enzyme has room for improvement in view of the industrial production of crystal 1-kestose.

[0006] As a next step, some of the inventors have successfully screened new enzymes having more favorable characteristics from Penicillium roqueforti and Scopulariopsis brevicaulis. These enzymes were able to turn 47% and 55% of sucrose into 1-kestose, respectively, and 7% and 4% to nystose (Japanese Patent Application No. 77534/1996, and Japanese Patent Application No. 77539/1996). These enzymes are inferior in productivity and stability to the enzyme derived from Aspergillus niger, and have room for improvement in view of the industrial production of crystal 1-kestose.

[0007] Thus, some of the inventors had paid attention to the procedure of genetic engineering as a process for improving the productivity of the enzyme, isolated the gene encoding β -fructofuranosidase from Penicillium roqueforti and Scopulariopsis brevicaulis, respectively, and conducted the structure analysis (PCT/JP97/00757). As a result, the translation regions encoding 565 amino acids and 574 amino acids as a mature protein were respectively deduced in the β -fructofuranosidase genes from Penicillium roqueforti and Scopulariopsis brevicaulis and their expression products were shown to have β -fructofuranosidase activity, like the β -fructofuranosidase gene from Aspergillus niger (L.M. Boddy et al., Curr. Genet., 24, 60-66 (1993)).

50 Summary of the Invention

[0008] The inventors have now found that the addition of 38 and 39 amino acids to the C-terminal of the β -fructofuranosidase genes from Penicillium roqueforti and Scopulariopsis brevicaulis, which were previously found by some of the inventors, improves its activity.

55 [0009] Thus, an object of the present invention is to provide a novel β -fructofuranosidase and its gene.

[0010] The novel β -fructofuranosidase according to the present invention is a polypeptide comprising the amino acid sequence of SEQ ID No. 1 or 3 or a homologue thereof.

[0011] Furthermore, the gene according to the present invention is a DNA encoding the above polypeptide.

EP 1 022 332 A1

[0012] The amino acid sequence of SEQ ID No. 1 or 3 according to the present invention is constructed by adding 38 and 39 amino acids to the C-terminals of the β -fructofuranosidase genes from Penicillium roqueforti and Scopulariopsis brevicaulis, which were previously found by some of the inventors as described above. It has been found that an intron actually exists at the region of the β -fructofuranosidase gene, which was presumed to encode the C-terminal amino acids by some of the present inventors and that the β -fructofuranosidase genes further encode 38 and 39 amino acids of the C-terminal. Surprisingly, the β -fructofuranosidase activity was remarkably improved by adding these amino acids to the C-terminal, as compared with the protein to which these sequences are not added.

Brief Description of the Drawing

[0013]

Figures 1A, B, C and D show the construction of expression vector pYPEN02 in which a gene encoding the enzyme protein consisting of the amino acid sequence of SEQ ID No. 1 is introduced, and expression vector pYPEN01 in which a gene encoding the enzyme protein consisting of the amino acid sequence from 1 to 565 of amino acid sequence of SEQ ID No. 1 is introduced.

Figures 2A and B show the construction of expression vector pYSCOP02 in which a gene encoding the enzyme protein consisting of the amino acid sequence of SEQ ID No. 3 is introduced, and expression vector pYSCOP01 in which a gene encoding the enzyme protein consisting of the amino acid sequence from 1 to 574 of amino acid sequence of SEQ ID No. 3 is introduced.

Detailed Description of the Invention β -fructofuranosidase

[0014] The polypeptide according to the present invention comprises the amino acid sequence of SEQ ID No. 1 or 3. This polypeptide having the amino acid sequence of SEQ ID No. 1 or 3 has enzymatic activity as β -fructofuranosidase. The polypeptide according to the present invention involves a homologue of the amino acid sequence of SEQ ID No. 1 or 3 as shown in the sequence listing. The term "homologue" refers to an amino acid sequence in which one or more amino acids (for example, one to several amino acids) are inserted, substituted or deleted in, or added to either or both of the terminals of, the amino acid sequence of SEQ ID Nos. 1 and 3 while retaining β -fructofuranosidase activity. Such a homologue can be selected and produced by those skilled in the art without undue experiments by referring to the sequence of SEQ ID No. 1 or 3.

[0015] The β -fructofuranosidase having the amino acid sequence of SEQ ID Nos. 1 and 3 according to the present invention has high fructosyltransferase activity and efficiently produces fructooligosaccharides. Specifically, when a sucrose solution at a concentration of 30 wt% or more is used as a substrate for reaction, the fructosyltransferase activity of β -fructofuranosidase having the amino acid sequence of SEQ ID No. 1 is at least 4 times higher, and the fructosyltransferase activity of β -fructofuranosidase having the amino acid sequence of SEQ ID No. 3 is at least 7 times higher than hydrolytic activity. Furthermore, 50% or more of sucrose is converted to fructooligosaccharides in both cases.

 β -fructofuranosidase gene

[0016] The novel gene encoding β -fructofuranosidase according to the present invention comprises a DNA sequence encoding the amino acid sequence of SEQ ID Nos. 1 and 3 or a homologue thereof.

[0017] Generally, a nucleotide sequence which encodes the amino acid sequence of a given protein can be easily determined from the reference chart known as "codon table". A variety of nucleotide sequence are available from those encoding the amino acid sequence of SEQ ID No. 1 or 3. Therefore, the term "a nucleotide sequence encoding the amino acid sequence of SEQ ID No. 1 or 3" refers to the meaning including the nucleotide sequence of SEQ ID No. 2 or 4, as well as nucleotide sequences which consist of the same codons as above allowing for degeneracy and encode the amino acid sequence of SEQ ID No. 1 or 3.

[0018] A preferred embodiment of the present invention provides, as a preferred example of the novel gene according to the present invention, a DNA fragment comprising the nucleotide sequence of SEQ ID No. 2 or 4.

[0019] As described above, the present invention encompasses a homologue of the amino acid sequence of SEQ ID No. 1 or 3. Therefore, the DNA fragment according to the present invention involves a nucleotide sequence which encodes such a homologue.

[0020] As the nucleotide sequence of the DNA fragment according to the present invention is determined, the DNA fragment may be obtained according to the procedure for the synthesis of a nucleic acid.

[0021] This sequence can also be obtained from Penicillium roqueforti or Scopulariopsis brevicaulis, preferably

EP 1 022 332 A1

Penicillium roqueforti IAM7254 or Scopulariopsis brevicaulis IFO4843, according to the procedure of genetic engineering.

Expression of β -fructofuranosidase Gene

[0022] The β -fructofuranosidase according to the present invention can be produced in a host cell which has been transformed by a DNA fragment encoding the enzyme. More specifically, a DNA fragment encoding the β -fructofuranosidase according to the present invention is introduced in a host cell in the form of a DNA molecule which is replicatable in the host cell and can express the above gene, particularly an expression vector, in order to transform the host cell. Then, the obtained transformant is cultivated.

[0023] Therefore, the present invention provides a DNA molecule which comprises a gene encoding the β -fructofuranosidase according to the present invention, particularly an expression vector. This DNA molecule is obtained by introducing a DNA fragment encoding the β -fructofuranosidase according to the present invention in a vector molecule. According to a preferred embodiment of the present invention, the vector is a plasmid.

[0024] The DNA molecule according to the present invention may be prepared by the standard technique of genetic engineering.

[0025] The vector applicable in the present invention can be selected as appropriate from viruses, plasmids, cosmid vectors, etc., considering the type of the host cell used. For example, a bacteriophage in the λ phage group or a plasmid in the pBR or pUC group may be used for E. coli host cells, a plasmid in the pUB group for Bacillus subtilis, and a vector in the YEp or YCp group for yeast.

[0026] It is preferable that the plasmid contain a selectable marker to ensure the selection of the obtained transformance, such as a drug-resistance marker or marker gene complementing an auxotrophic mutation. Preferred example of marker genes include ampicillin-resistance gene, kanamycin-resistance gene, and tetracycline-resistance gene for bacterium host cells; N-(5'-phosphoribosyl)-anthranilate isomerase gene (TRP1), orotidine-5'-phosphate decarboxylase gene (URA3), and β -isopropylmalate dehydrogenase gene (LEU2) for yeast; and hygromycin-resistance gene (hph), bialaphos-resistance gene (bar), and nitrate reductase gene (niaD) for mold.

[0027] It is also preferable that the DNA molecule for use as an expression vector according to the present invention contain nucleotide sequence necessary for the expression of the β -fructofuranosidase gene, including transcription and translation control signals, such as a promoter; a transcription initiation signal, a ribosome binding site, a translation termination signal, and a transcription termination signal.

[0028] Examples of preferred promoters include, in addition to the promoter on the inserted fragment which is able to function in the host, promoters such as those of lactose operon (lac), and tryptophan operon (trp) for E. coli; promoters such as those of alcohol dehydrogenase gene (ADH), acid phosphatase gene (PHO), galactose regulated gene (GAL), and glyceraldehyde-3-phosphate dehydrogenase gene (GPD) for yeast; and promoters such as those of α -amylase gene (amy) and cellobiohydrolase I gene (CBHI) for mold.

[0029] When the host cell is Bacillus subtilis, yeast or mold, it is also advantageous to use a secretion vector to allow it to extracellularly secrete the produced recombinant β -fructofuranosidase. Any host cell with an established host-vector system may be used, preferably yeast, mold, etc. It is preferable also to use the mold fungus having no β -fructofuranosidase activity described in PCT/JP97/00757.

[0030] A novel recombinant enzyme produced by the transformant described above is obtained by the following procedure: first, the host cell described above is cultivated under suitable conditions to obtain the supernatant or cell bodies from the resultant culture, using a known technique such as centrifugation; cell bodies should be further suspended in a suitable buffer solution, then homogenized by freeze-and-thaw, ultrasonic treatment, or mortar, followed by centrifugation or filtration to separate a cell body extract containing the novel recombinant enzyme.

[0031] The enzyme can be purified by combining the standard techniques for separation and purification. Examples of such as techniques include processes such as heat treatment, which rely on the difference in thermal resistance; processes such as salt sedimentation and solvent sedimentation, which rely on the difference in solubility; processes such as dialysis, ultrafiltration and gel filtration, and SDS-polyacrylamide gel electrophoresis, which rely on the difference in molecular weight; processes such as ion exchange chromatography, which rely on the difference in electric charge; processes such as affinity chromatography, which rely on specific affinity; processes such as hydrophobic chromatography and reversed-phase partition chromatography, which rely on the difference in hydrophobicity; and processes such as isoelectric focusing, which rely on the difference in isoelectric point.

Production of fructooligosaccharides using the β -fructofuranosidase

[0032] The present invention further provides a process for producing fructooligosaccharide using the recombinant host or recombinant β -fructofuranosidase described above.

[0033] In the process for producing fructooligosaccharides according to the present invention, the recombinant host

EP 1 022 332 A1

or recombinant β -fructofuranosidase described above is brought into contact with sucrose.

[0034] The mode and conditions where the recombinant host or recombinant β -fructofuranosidase according to the present invention comes in contact with sucrose are not limited in any way provided that the novel recombinant enzyme is able to act on sucrose. A preferred embodiment for contact in solution is as follows: The sucrose concentration may be selected as appropriate in the range where sucrose can be dissolved. However, considering the conditions such as the specific activity of the enzyme and reaction temperature, the concentration should generally fall in the range of 5% to 80%, preferably 30% to 70%. The temperature and pH for the reaction of sucrose by the enzyme should preferably be optimized for the characteristics of the novel recombinant enzyme. Therefore, the reasonable conditions are about 30°C to 80°C, pH 4 to 10, preferably 40°C to 70°C, pH 5 to 7.

[0035] The degree of purification of the novel recombinant enzyme may be selected as appropriate. The enzyme may be used either as unpurified in the form of supernatant from a transformant culture or cell body homogenate, as purified after processed in various purification steps, or as isolated after processed by various purification means.

[0036] Furthermore, the enzyme may be brought into contact with sucrose as fixed on a carrier using the standard technique.

[0037] The fructooligosaccharides thus produced are purified from the resulting solution according to known procedures. For example, the solution may be heated to inactivate the enzyme, decolorized using activated carbon, then desalted using ion exchange resin.

Examples

Example 1: Determination of translation region of β -fructofuranosidase gene from Penicillium roqueforti IAM7254

[0038] A DNA fragment of about 2 kbp containing the β -fructofuranosidase gene from Aspergillus niger was amplified by PCR using a chromosomal DNA prepared from Aspergillus niger ATCC20611 according to the standard procedure as a template and synthetic DNAs of SEQ ID Nos. 5 and 6 as primers. This DNA fragment was fractionated by agarose gel electrophoresis, extracted according to the standard procedure, purified, and then dissolved in sterilized water to 0.1 $\mu\text{g}/\mu\text{l}$ to prepare a DNA sample for probe.

[0039] In the next step, a chromosomal DNA from Penicillium roqueforti IAM7254 was prepared, about 20 μg of the chromosomal DNA was digested completely with EcoRI, followed by agarose gel electrophoresis to recover about 4 kbp DNA fragments.

[0040] The recovered DNA fragments of about 4 kbp (about 0.5 μg) were ligated with 1 μg of $\lambda\text{gt}10$ vector which had been digested with EcoRI and treated with phosphatase, packaged using an *in vitro* packaging kit, GIGAPACK II Gold (Stratagene L.L.C.), and then introduced in E. coli NM514, to prepare a library.

[0041] A probe was prepared from DNA sample for probe above described. As a result of plaque hybridization using ECL Direct DNA/RNA Labelling & Detection System (Amersham International), 4 clones turned out positive in about 25,000 plaques. These positive clones were purified by a second screening to prepare phage DNA, which was then analyzed using restriction enzymes. The result showed that all the clones had an identical EcoRI fragment of about 4 kbp.

[0042] The EcoRI fragments of about 4 kbp were subdivided into a small fragment to select the desired DNA region using restriction enzymes, then subcloned to plasmid vector pUC118 or pUC119. The plasmid DNA was obtained from the subclone according to the standard procedure and sequenced using ALFred DNA Sequencer (Pharmacia) as shown in SEQ ID No.7.

[0043] The sequence consisting 50 bases from 1695 to 1744 in this sequence was identified as an intron because it showed a typical intron structure of filamentous fungi. As a result, the sequence of SEQ ID No. 2 as a sequence encoding protein was obtained by deleting the intron from the sequence of SEQ ID No. 7. The encoded amino acid sequence was shown in SEQ ID No. 1.

Example 2: Expression of β -fructofuranosidase gene from Penicillium roqueforti IAM7254 in Saccharomyces cerevisiae

[0044] Plasmid pYPEN01 and pYPEN02 for expressing the β -fructofuranosidase gene from Penicillium roqueforti were prepared as follows (Figure 1A, B, C and D).

[0045] pYPR2831 (H. Horiuchi et al., Agric. Biol. Chem., 54, 1771-1779, 1990) was digested with EcoRI and Sall, and then its terminals were blunted with T4 DNA polymerase. The obtained fragment was ligated with BamHI linker (5'-CGGATCCG-3'), digested with BamHI, followed by self-ligation to obtain vector pY2831 for expression in yeast.

[0046] Next, single-stranded DNA was prepared from the plasmid pPRS01 obtained by inserting an about 4 kbp EcoRI DNA fragment containing the β -fructofuranosidase gene prepared in Example 1 into plasmid pUC118. Using the single-stranded DNA as a template and a synthetic DNA of SEQ ID No. 8 as a primer, the translated region of the β -fructofuranosidase gene was subjected to site-specific mutagenesis to disrupt the BamHI site without changing the

EP 1 022 332 A1

encoded amino acid sequence (pPRS02).

[0047] A part of the translated region of the β -fructofuranosidase gene was prepared as an about 1.8 kbp BamHI fragment by PCR using plasmid pPRS02 as a template and synthetic DNAs of SEQ ID Nos. 9 and 10 as primers, and inserted into the BamHI site of plasmid pY2831 to prepare pYPEN01. Thus, plasmid pYPEN01 is designed to produce an enzyme protein comprising an amino acid sequence from 1 to 565 in the amino acid sequence of SEQ ID No. 1, which is a mature β -fructofuranosidase following secretion signal sequence.

[0048] Further, a DNA fragment containing the translated region of the β -fructofuranosidase gene was prepared as an about 1.8 kbp BamHI fragment by PCR using plasmid pPRS02 as a template and synthetic DNAs of SEQ ID Nos. 9 and 11 as primers, and inserted into the BamHI site of plasmid pUC118 to prepare plasmid pPRS03. A single-stranded DNA was prepared from plasmid pPRS03. As a result of site-specific mutagenesis using this as a template and a synthetic DNA of SEQ ID No. 12 as a primer, the intron sequence was removed (pPRS04). The translated region of the β -fructofuranosidase gene was prepared as an about 1.8 kbp BamHI fragment from plasmid pPRS04, and inserted into the BamHI site of plasmid pY2831 to prepare plasmid pYPEN02. Thus, plasmid pYPEN02 is designed to produce an enzyme protein comprising an amino acid sequence of SEQ ID No. 1, which is a mature β -fructofuranosidase following secretion signal sequence.

[0049] Plasmids pYPEN01 and pYPEN02 were introduced into Saccharomyces cerevisiae MS-161 (Suc⁻, ura3, trp1) by the lithium-acetate method (Ito, H. et al., J. Bacteriol., 153, 163-168, 1983) to obtain transformants. The transformants were cultivated in an SD-Ura medium (0.67% yeast nitrogen base (Difco), 2% glucose and 50 μ g/ml uracil) at 30°C overnight. The culture was seeded in a production medium (0.67% yeast nitrogen base Difco, 2% glucose, 2% casamino acid and 50 μ g/ml uracil) at a final concentration of 1% and cultivated at 30°C for 2 days. The culture supernatant was analyzed for β -fructofuranosidase activity, in units, i.e., the quantity of free glucose (μ mol) released in 1 minute in 10 wt% sucrose solution, pH 5.5, at 40°C for 60 minutes. As a result, the transformant with plasmid pYPEN01 exhibited 4×10^{-4} units/ml or less of activity while the transformant with plasmid pYPEN02 exhibited 0.38 units/ml of activity.

Example 3: Determination of the translated region of β -fructofuranosidase gene from Scopulariopsis brevicaulis IFO4843

[0050] The chromosomal DNA was prepared from Scopulariopsis brevicaulis IFO4843. About 20 μ g of a chromosomal DNA sample was completely digested with EcoRI, and electrophoresed through an agarose gel to recover an about 10 kbp DNA fragment.

[0051] The recovered DNA fragment of about 10 kbp (about 0.5 μ g) were ligated with 1 μ g of λ DASHII vector digested with HindIII and EcoRI, and packaged using an *in vitro* packaging kit, GIGAPACK II Gold (Stratagene L.L.C.), then introduced in E. coli XL1-Blue MRA (P2), to prepare a library.

[0052] As a result of plaque hybridization using ECL Direct DNA/RNA Labelling & Detection System (Amersham International), with the about 2 kbp DNA fragment used in Example 1 as a probe, 3 clones turned out positive in about 15,000 plaques. These positive clones were purified by a second screening to prepare phage DNA, which was then analyzed using restriction enzymes. The result showed that all the clones had an identical EcoRI fragment of about 10 kbp.

[0053] These EcoRI fragments of about 10 kbp were subdivided into a small fragment to select the desired DNA region using restriction enzymes, then subcloned to plasmid vector pUC118 or pUC119. The plasmid DNA was obtained from the subclone according to the standard procedure and sequenced using ALFred DNA Sequencer (Pharmacia) as shown in SEQ ID No.13.

[0054] The sequence comprising 55 bases from 1722 to 1776 in this sequence was identified as an intron because it showed a typical intron structure of filamentous fungi. As a result, the sequence of SEQ ID No. 4 as a sequence encoding protein was obtained by deleting the intron from the sequence of SEQ ID No. 13. The encoded amino acid sequence was shown SEQ ID No. 3.

Example 4: Expression of β -fructofuranosidase gene from Scopulariopsis brevicaulis IFO4843 in Saccharomyces cerevisiae

[0055] Plasmid pYSCOP01 and pYSCOP02 for expressing the β -fructofuranosidase gene from Scopulariopsis brevicaulis were prepared as follows (Figure 2A and B).

[0056] A part of the translated region of the β -fructofuranosidase gene was prepared as an about 1.8 kbp BamHI fragment by PCR using about 10 kbp EcoRI DNA fragment prepared in Example 3 containing the β -fructofuranosidase gene as a template and synthetic DNAs of SEQ ID Nos. 14 and 15 as primers, and inserted into the BamHI site of plasmid pY2831 to prepare pYSCOP01. Thus, plasmid pYSCOP01 is designed to produce an enzyme protein comprising an amino acid sequence from 1 to 574 in the amino acid sequence of SEQ ID No. 3, which is a mature β -fructofuranosi-

EP 1 022 332 A1

dase following secretion signal sequence.

[0057] Next, a DNA fragment containing the translated region of the β -fructofuranosidase gene was prepared as an about 1.9 kbp BamHI fragment by PCR using an about 10 kbp EcoRI fragment containing the β -fructofuranosidase gene as a template and synthetic DNAs of SEQ ID Nos. 14 and 16 as primers, and inserted into the BamHI site of plasmid pUC118 to prepare plasmid pSCB01. A single-stranded DNA was prepared from plasmid pSCB01. As a result of site-specific mutagenesis using this as a template and the synthetic DNA of SEQ ID No. 17 as a primer, the intron sequence was removed (pSCB02). The translated region of the β -fructofuranosidase gene was prepared as an about 1.9 kbp BamHI fragment from plasmid pSCB02, and inserted into the BamHI site of plasmid pY2831 to prepare plasmid pYSCOP02. Thus, plasmid pYSCOP02 is designed to produce an enzyme protein comprising an amino acid sequence of SEQ ID No. 3, which is a mature β -fructofuranosidase following secretion signal sequence.

[0058] Plasmids pYSCOP01 and pYSCOP02 were introduced into Saccharomyces cerevisiae MS-161 (Suc⁺, ura3, trp1) by the lithium-acetate method to obtain transformants. The transformants were cultivated in an SD-Ura medium at 30°C overnight. The culture was seeded a production medium at a final concentration of 1% and cultivated at 30°C for 2 days. The culture supernatant was analyzed for β -fructofuranosidase activity in the same manner as described in Example 2. As a result, the transformant with plasmid pYSCOP01 exhibited 4×10^{-4} units/ml or less of activity, while the transformant with plasmid pYSCOP02 exhibited 6.5×10^{-3} units/ml of activity.

EP 1 022 332 A1

SEQUENCE LISTING

5 <110> MEIJI SEIKA KAISHA, LTD.

10 <120> Beta-fructofuranosidase and its gene

<130> 116875-475

15 <140>

<141> 1998-9-10

20 <150> JP/245154/1997

<151> 1997-9-10

25 <160> 17

30 <170> PatentIn Ver. 2.0

35 <210> 1

<211> 603

<212> PRT

40 <213> Penicillium roqueforti IAM7254

<220>

<221> mat peptide

45 <222> (1)... (603)

<400> 1

50 Val Asp Phe His Thr Pro Ile Asp Tyr Asn Ser Ala Pro Pro Asn Leu

1 5 10 15

55

EP 1 022 332 A1

Ser Thr Leu Ala Asn Ala Ser Leu Phe Lys Thr Trp Arg Pro Arg Ala

20

25

30

5

His Leu Leu Pro Pro Ser Gly Asn Ile Gly Asp Pro Cys Gly His Tyr

35

40

45

10

Thr Asp Pro Lys Thr Gly Leu Phe His Val Gly Trp Leu Tyr Ser Gly

50

55

60

Ile Ser Gly Ala Thr Thr Asp Asp Leu Val Thr Tyr Lys Asp Leu Asn

15

65

70

75

80

Pro Asp Gly Ala Pro Ser Ile Val Ala Gly Gly Lys Asn Asp Pro Leu

85

90

95

20

Ser Val Phe Asp Gly Ser Val Ile Pro Ser Gly Ile Asp Gly Met Pro

100

105

110

Thr Leu Leu Tyr Thr Ser Val Ser Tyr Leu Pro Ile His Trp Ser Ile

25

115

120

125

Pro Tyr Thr Arg Gly Ser Glu Thr Gln Ser Leu Ala Val Ser Tyr Asp

130

135

140

30

Gly Gly His Asn Phe Thr Lys Leu Asn Gln Gly Pro Val Ile Pro Thr

145

150

155

160

Pro Pro Phe Ala Leu Asn Val Thr Ala Phe Arg Asp Pro Tyr Val Phe

35

165

170

175

Gln Ser Pro Ile Leu Asp Lys Ser Val Asn Ser Thr Gln Gly Thr Trp

40

180

185

190

Tyr Val Ala Ile Ser Gly Gly Val His Gly Val Gly Pro Cys Gln Phe

195

200

205

45

Leu Tyr Arg Gln Asn Asp Ala Asp Phe Gln Tyr Trp Glu Tyr Leu Gly

210

215

220

Gln Trp Trp Lys Glu Pro Leu Asn Thr Thr Trp Gly Lys Gly Asp Trp

50

225

230

235

240

Ala Gly Gly Trp Gly Phe Asn Phe Glu Val Gly Asn Val Phe Ser Leu

55

EP 1 022 332 A1

	245	250	255
5	Asn Ala Glu Gly Tyr Ser Glu Asp Gly Glu Ile Phe Ile Thr Leu Gly		
	260	265	270
	Ala Glu Gly Ser Gly Leu Pro Ile Val Pro Gln Val Ser Ser Ile Arg		
10	275	280	285
	Asp Met Leu Trp Val Thr Gly Asn Val Thr Asn Asp Gly Ser Val Thr		
	290	295	300
15	Phe Lys Pro Thr Met Ala Gly Val Leu Asp Trp Gly Val Ser Ala Tyr		
	305	310	315
	Ala Ala Ala Gly Lys Ile Leu Pro Ala Ser Ser Gln Ala Ser Thr Lys		320
20	325	330	335
	Ser Gly Ala Pro Asp Arg Phe Ile Ser Tyr Val Trp Leu Thr Gly Asp		
	340	345	350
25	Leu Phe Glu Gln Val Lys Gly Phe Pro Thr Ala Gln Gln Asn Trp Thr		
	355	360	365
	Gly Ala Leu Leu Leu Pro Arg Glu Leu Asn Val Arg Thr Ile Ser Asn		
30	370	375	380
	Val Val Asp Asn Glu Leu Ser Arg Glu Ser Leu Thr Ser Trp Arg Val		
	385	390	395
35	Ala Arg Glu Asp Ser Gly Gln Ile Asp Leu Glu Thr Met Gly Ile Ser		
	405	410	415
	Ile Ser Arg Glu Thr Tyr Ser Ala Leu Thr Ser Gly Ser Ser Phe Val		
40	420	425	430
	Glu Ser Gly Lys Thr Leu Ser Asn Ala Gly Ala Val Pro Phe Asn Thr		
	435	440	445
45	Ser Pro Ser Ser Lys Phe Phe Val Leu Thr Ala Asn Ile Ser Phe Pro		
	450	455	460
50	Thr Ser Ala Arg Asp Ser Gly Ile Gln Ala Gly Phe Gln Val Leu Ser		
	465	470	475
			480
55			

EP 1 022 332 A1

Ser Ser Leu Glu Ser Thr Thr Ile Tyr Tyr Gln Phe Ser Asn Glu Ser

485

490

495

5

Ile Ile Val Asp Arg Ser Asn Thr Ser Ala Ala Ala Arg Thr Thr Ala

500

505

510

10

Gly Ile Leu Ser Asp Asn Glu Ala Gly Arg Leu Arg Leu Phe Asp Val

515

520

525

Leu Arg Asn Gly Lys Glu Gln Val Glu Thr Leu Glu Leu Thr Ile Val

15

530

535

540

Val Asp Asn Ser Val Leu Glu Val Tyr Ala Asn Gly Arg Phe Ala Leu

545

550

555

560

20

Gly Thr Trp Ala Arg Ser Trp Tyr Ala Asn Ser Thr Lys Ile Asn Phe

565

570

575

Phe His Asn Gly Val Gly Glu Ala Thr Phe Glu Asp Val Thr Val Phe

25

580

585

590

Glu Gly Leu Tyr Asp Ala Trp Pro Gln Arg Lys

595

600

30

<210> 2

<211> 1809

35

<212> DNA

<213> Penicillium roqueforti IAM7254

<220>

40

<221> mat peptide

<222> (1)... (1809)

45

<400> 2

gttgatttcc ataccccgat tgactataac tcggctccgc caaaccttcc taccctggca 60

50

aacgcacatc ttccaagac atggagaccc agagcccatc ttctccctcc atctgggaac 120

ataggcgacc cgtgcgggca ctataccgat cccaagactg gtctcttcca cgtgggttgg 180

55

EP 1 022 332 A1

	ctttacagtg ggatttcggg agcgacaacc gacgatctcg ttacctataa agacctcaat	240
5	cccgatggag ccccgtaaat tgttgcagga ggaaagaacg accctctttc tgtcttcgat	300
	ggctcgggta ttccaagcgg tatagacggc atgccaactc ttctgtatac ctctgtatca	360
	tacctcccaa tccactggtc catcccctac acccggggaa gcgagacaca atccttggcc	420
10	gtttcctatg acggiggtca caacttcacc aagctcaacc aaggggccgt gatccctacg	480
	cttccgtttg ctctcaatgt caccgccttc cgtgacccct acgttttcca aagcccaatt	540
	ctggacaaat ctgtcaatag tacccaagga acatgggatg tcgccatata tggcgggtgc	600
15	cacgggtgctg gaccttgta gttectctac cgtcagaacg acgcagattt tcaatatgg	660
	gaatatctcg ggcaatgggtg gaaggagccc cttaatacca ctgggggaaa gggtagctgg	720
	gccgggggtt ggggcctcaa ctttgagggt ggcaacgtct ttagtctgaa tgcagagggg	780
20	tatagtgaag acggcgagat attcataacc ctcggtgctg agggttcggg acttcccatc	840
	gttccctcaag tctcctctat tcgcgatatg ctgtgggtga ccggcaatgt cacaaatgac	900
	ggctctgtca ctttcaagcc aaccaatggcg ggtgtgcttg actggggcgt gtcggcatat	960
25	gttgcctcag gcaagatctt gccggccagc tctcaggcat ccacaaagag cggtgccccc	1020
	gatcgggtca tttcctatgt ctggctcact ggagatctat tcgagcaagt gaaaggattc	1080
30	cttaccgctc aacaaaactg gaccggggcc ctcttactgc ccgagagct gaatgtccgc	1140
	actatctcta acgtgggtga taacgaactt tcgcgtgagt ccttgacatc gtggcgctg	1200
	gcccgcgaag actcgggtca gatcgacctt gaaacaatgg gaatctcaat tccaggzag	1260
35	acttacagcg ctctcaccac cggctcatct ttgttcgagt ctggtaaaac gttgtcgzat	1320
	gctggagcag tgccttcaaa tacctcacc tcaagcaagt tcttcgtgct gacagcaaat	1380
	atatctttcc cgacctctgc ccgtgactct ggcatccagg ctggtttcca ggttttatcc	1440
40	tctagtcctg agtctacaac tatctactac caattctcca acgagttcat catcgtcgac	1500
	cgcagcaaca cgagtgcctg ggcgagaaca actgctggga tcttcagtga taacgaggcg	1560
45	ggacgtctgc gccctctcga cgtgttgcga aatggaaaag aacaggttga aactttggag	1620
	ctcactatcg tggtagataa tagtgfactg gaagtatatg ccaatggacg ctttgctcta	1680
	ggcacttggg ctcggtcttg gtacgccaac tcgactaaaa ttaacttctt ccataacggc	1740
50	gtgggagaag cgacattcga agatgtgacg gtctttgaag gactgtatga tgcctggcca	1800
	caaaggaag	1809

55

EP 1 022 332 A1

<210> 3

<211> 613

<212> PRT

<213> Scopulariopsis brevicaulis IF04843

<220>

<221> mat peptide

<222> (1)... (613)

<400> 3

Gln Pro Thr Ser Leu Ser Ile Asp Asn Ser Thr Tyr Pro Ser Ile Asp

1 5 10 15

Tyr Asn Ser Ala Pro Pro Asn Leu Ser Thr Leu Ala Asn Asn Ser Leu

20 25 30

Phe Glu Thr Trp Arg Pro Arg Ala His Val Leu Pro Pro Gln Asn Gln

35 40 45

Ile Gly Asp Pro Cys Met His Tyr Thr Asp Pro Glu Thr Gly Ile Phe

50 55 60

His Val Gly Trp Leu Tyr Asn Gly Asn Gly Ala Ser Gly Ala Thr Thr

65 70 75 80

Glu Asp Leu Val Thr Tyr Gln Asp Leu Asn Pro Asp Gly Ala Gln Met

85 90 95

Ile Leu Pro Gly Gly Val Asn Asp Pro Ile Ala Val Phe Asp Gly Ala

100 105 110

Val Ile Pro Ser Gly Ile Asp Gly Lys Pro Thr Met Met Tyr Thr Ser

115 120 125

Val Ser Tyr Met Pro Ile Ser Trp Ser Ile Ala Tyr Thr Arg Gly Ser

130 135 140

Glu Thr His Ser Leu Ala Val Ser Ser Asp Gly Gly Lys Asn Phe Thr

145 150 155 160

EP 1 022 332 A1

Lys Leu Val Gln Gly Pro Val Ile Pro Ser Pro Pro Phe Gly Ala Asn

		165		170		175
5	Val Thr Ser Trp Arg Asp Pro Phe Leu Phe Gln Asn Pro Gln Phe Asp					
		180		185		190
10	Ser Leu Leu Glu Ser Glu Asn Gly Thr Trp Tyr Thr Val Ile Ser Gly					
		195		200		205
	Gly Ile His Gly Asp Gly Pro Ser Ala Phe Leu Tyr Arg Gln His Asp					
15		210		215		220
	Pro Asp Phe Gln Tyr Trp Glu Tyr Leu Gly Pro Trp Trp Asn Glu Glu					
		225		230		235
20	Gly Asn Ser Thr Trp Gly Ser Gly Asp Trp Ala Gly Arg Trp Gly Tyr					
		245		250		255
	Asn Phe Glu Val Ile Asn Ile Val Gly Leu Asp Asp Asp Gly Tyr Asn					
25		260		265		270
	Pro Asp Gly Glu Ile Phe Ala Thr Val Gly Thr Glu Trp Ser Phe Asp					
		275		280		285
30	Pro Ile Lys Pro Gln Ala Ser Asp Asn Arg Glu Met Leu Trp Ala Ala					
		290		295		300
	Gly Asn Met Thr Leu Glu Asp Gly Asp Ile Lys Phe Thr Pro Ser Met					
35		305		310		315
						320
	Ala Gly Tyr Leu Asp Trp Gly Leu Ser Ala Tyr Ala Ala Ala Gly Lys					
		325		330		335
40	Glu Leu Pro Ala Ser Ser Lys Pro Ser Gln Lys Ser Gly Ala Pro Asp					
		340		345		350
45	Arg Phe Val Ser Tyr Leu Trp Leu Thr Gly Asp Tyr Phe Glu Gly His					
		355		360		365
	Asp Phe Pro Thr Pro Gln Gln Asn Trp Thr Gly Ser Leu Leu Leu Pro					
50		370		375		380
	Arg Glu Leu Ser Val Gly Thr Ile Pro Asn Val Val Asp Asn Glu Leu					

55

EP 1 022 332 A1

	385	390	395	400
5	Ala Arg Glu Thr Gly Ser Trp Arg Val Gly Thr Asn Asp Thr Gly Val			
		405	410	415
	Leu Glu Leu Val Thr Leu Lys Gln Glu Ile Ala Arg Glu Thr Leu Ala			
10		420	425	430
	Glu Met Thr Ser Gly Asn Ser Phe Thr Glu Ala Ser Arg Asn Val Ser			
	435	440	445	
15	Ser Pro Gly Ser Thr Ala Phe Gln Gln Ser Leu Asp Ser Lys Phe Phe			
	450	455	460	
	Val Leu Thr Ala Ser Leu Ser Phe Pro Ser Ser Ala Arg Asp Ser Asp			
20	465	470	475	480
	Leu Lys Ala Gly Phe Glu Ile Leu Ser Ser Glu Phe Glu Ser Thr Thr			
		485	490	495
25	Val Tyr Tyr Gln Phe Ser Asn Glu Ser Ile Ile Ile Asp Arg Ser Asn			
	500	505	510	
	Ser Ser Ala Ala Ala Leu Thr Thr Asp Gly Ile Asp Thr Arg Asn Glu			
30	515	520	525	
	Phe Gly Lys Met Arg Leu Phe Asp Val Val Glu Gly Asp Gln Glu Arg			
35	530	535	540	
	Ile Glu Thr Leu Asp Leu Thr Ile Val Val Asp Asn Ser Ile Val Glu			
	545	550	555	560
40	Val His Ala Asn Gly Arg Phe Ala Leu Ser Thr Trp Val Arg Ser Trp			
		565	570	575
	Tyr Glu Ser Ser Lys Asp Ile Lys Phe Phe His Asp Gly Asp Ser Thr			
45	580	585	590	
	Val Gln Phe Ser Asn Ile Thr Val Tyr Glu Gly Leu Phe Asp Ala Trp			
	595	600	605	
50	Pro Glu Arg Ala Arg			
	610			

55

EP 1 022 332 A1

<210> 4

5

<211> 1839

<212> DNA

<213> Scopulariopsis brevicaulis IF04843

10

<220>

<221> mat peptide

<222> (1)... (1839)

15

<400> 4

20

caacctacgt ctctgtcaat cgacaattcc acgtatccct ctatcgacta caactccgcc 60

cctccaaacc tctcgactct tgccaacaac agcctcttcg agacatggag gccgagggca 120

cacgtccctc cgccccagaa ccagatcggc gatccgltga tgcactacac cgaccccgag 180

25

acaggaatct tccacgtcgg ctggctgtac aacggcaatg gcgcttcgg cgccacgacc 240

gaggatctcg tcacctatca ggatctcaac cccgacggag cgcagatgat ccttccgggt 300

ggigtgaatg accccattgc tgtctttgac ggcgcggtta ttcccagtgg cattgatggg 360

30

aaaccaccca tgatgtatag ctcggtgtca tacatgccca tctcctggag catcgcttac 420

accaggggaa gcgagaccca ctctctcgca gtgtcttcg acggcggtia gaacttcacc 480

aagctggltc agggccccgt cattctctcg cctcccttcg gcgccaacgt gaccagctgg 540

35

cgtgacccct tctgtttcca aaacccccag ttcgactctc tcttcgaaag cgagaacggc 600

acgtgggtaca ccgttatctc tggltggcatc cacggtgacg gccccctccg gttctctac 660

40

cgtcagcacg accccgactt ccagttactg gagttacclt gaccgtgggt gaacgaggaa 720

gggaactcga cctggggcag cggtgactgg gctggccgggt ggggctacaa cttcgaggtc 780

atcaacattg tgggtcttga cgatgatggc tacaaccccg acggtgaaat ctttgccacg 840

45

gtaggtaccg aatggtcgtt tgaccccatc aaaccgcagg cctcggacaa cagggagatg 900

ctctgggccc cgggcaacat gactctcgag gacggcgata tcaagttcac gccaaagcatg 960

gcgggctacc tgcactgggg tctatcggcg tatgccgccc ctggcaagga gctgccccgt 1020

50

tcttcaaagc cttcgacagaa gagcggltcg ccggaccgggt tctgtctgta cctgtggctc 1080

accggtgact acttcgaggg ccacgacttc cccaccccg c agcagaattg gaccggctcg 1140

55

EP 1 022 332 A1

cttttgccttc cgcgtgagct gagcgtcggg acgattccca acgttgtcga caacgagctt 1200

5

gctcgcgaga cgggctcttg gagggttggc accaacgaca ctggcgigct tgagctggtc 1260

actctgaagc aggagattgc tcgcgagacg ctggctgaaa tgaccagcgg caactccttc 1320

accgaggcga gcaggaatgt cagctcgccc ggatctaccg ccttcagca gtccctggat 1380

10

tccaagtctt tcgtcctgac cgcctcgctc tcttccctt cgtcggctcg cgactccgac 1440

ctcaaggctg gtttcgagat cctgtcgtcc gagtttgagt cgaccacggt ctactaccag 1500

ttttccaacg agtccatcat catlgaccgg agcaactcga gtgcigccgc ctlgactacc 1560

15

gatggaatcg acaccgcga cgagtttggc aagatgcgcc tgtttgatgt tgtcgagggt 1620

gaccaggagc gtatcgagac gctcgatcic actattgttg ttgataactc gatcgttgag 1680

gttcatgcca acgggcgatt cgcctcgagc acttgggttc gttcgttgga cgagtcgtcc 1740

20

aaggacatca agttcttcca cgalggcgac agcacggttc agttctcgaa catcacgctc 1800

tacgagggac tgtttgacgc ctggccggag cgggccagg 1839

25

<210> 5

<211> 20

30

<212> DNA

<213> Artificial Sequence

35

<400> 5

caatgaagct caccactacc

40

<210> 6

<211> 20

<212> DNA

45

<213> Artificial Sequence

50

<400> 6

atcccggtea atttctctcc

55

EP 1 022 332 A1

<210> 7

<211> 1809

<212> DNA

<213> *Penicillium roqueforti* IAM7254

<220>

<221> CDS

<222> (1)... (1694)

<221> intron

<222> (1695)... (1744)

<221> CDS

<222> (1745)... (1859)

<400> 1

gttagatttcc ataccaccgat tgactataac tcggctccgc caaacctttc taccctggca 60
 aacgcattctc ttttcaagac atggagaccc agagcccatc ttctccctcc atctgggaac 120
 ataggcgacc cgtgcgggca ctataccgat cccaagactg gtctcttcca cgtgggttgg 180
 ctttacagtg ggatttcggg agegacaacc gacgatctcg ttacctataa agacctcaat 240
 cccgatggag ccccgicaat tgttgcagga gzaaagaacg accctctttc tgicttccat 300
 ggctcgggtca ttccaagcgg tatagacggc atgccaactc ttctgtatac ctctgtatca 360
 tacctcccaa tccactgggc catcccttac acccggggaa gcgagacaca atccttggcc 420
 gtttcctatg acggtgggtca caacttcacc aagctcaacc aagggcccggt gatccctacg 480
 cctccgtttg ctctcaatgt caccgctttc cgtgacccct acgttttcca aagcccaatt 540
 ctggacaaat ctgtcaatag tacccaagga acatggtaig tcgccatatac tggcgggtgc 600
 cacgggtgtcg gaccttggtc gttcctctac cgtcagaacg acgcagattt tcaatatagg 660
 gaatatctcg ggcaatgggt gaaggagccc cttaatacca ctigggggaaa gggtagctgg 720
 gccggggggtt ggggccttcaa ctttgagggt ggcaacgtct ttagtctgaa tgcagagggg 780
 tatagtgaag acggcgagat attcataacc ctcggtgtcg agggttcggg acttcccatc 840
 gttcctcaag tctcctctat tcgcgataig ctgtgggtga ccggcaatgt cacaaatgac 900

EP 1 022 332 A1

ggctctgtca ctttcaagcc aaccaaggcg ggtgtgttgc actggggcgt gtcggcatat 960

5

gtgtgtgcag gcaagatcct gccggccagc tctcaggcat ccacaaagag cggtagccccc 1020

gatacgttca tttcctatgt ctggctcact ggagatctat tcgagcaagt gaaaggattc 1080

cctaccgctc aacaaaacig gaccggggcc ctcttactgc cgcgagagct gaatgtccgc 1140

10

actatctcta acgtgttgga taacgaactt tcgcgtgagt ccttgacalc gtggcgcgtg 1200

gcccgcgaag actctgttca gatcgacctt gaaacaatgg gaatctcaat ttccaggagg 1260

acttacagcg ctctcacatc cggctcatct ttgttcgagt ctggtaaaac gttgtcgaat 1320

15

gctggagcag tgcccttcaa taccacacc tcaagcaagt tcttcgtgct gacagcaaat 1380

atatctttcc cgacctctgc ccgtgactct ggcatccagg ctggtttcca ggttttatcc 1440

tctagtcttg agtctacaac tatctactac caattctcca acgagtccat catcgtcgac 1500

20

cgcagcaaca cgagtgcigc ggcgagaaca actgctggga tcttcagtga taacgaggcg 1560

ggacgictgc gcctcttcga cgtgttgca aatggaaaag aacaggttga aacittggag 1620

ctcactatcg tggtagataa tagtgtactg gaagtatatg ccaatggacg ctttgtctta 1680

25

ggcacttggg ctcggttaagt ctctcttgtt taiggaagat tggtaaaaaa ctaaccgcat 1740

gaaggicttg gtacgccaac tcgactaaaa ttaacttctt ccataacggc gtgggagaag 1800

cgacattcga agatgtgacg gtctttgaag gacigtatga tgccctggcca caaaggaag 1859

30

<210> 8

<211> 22

35

<212> DNA

<213> Artificial Sequence

40

<400> 8

caactgctgg catcctcagt ga

45

<210> 9

<211> 29

50

<212> DNA

<213> Artificial Sequence

55

EP 1 022 332 A1

<400> 9

5 gcggatccat gaagctatca aatgcaatc

<210> 10

10

<211> 26

<212> DNA

<213> Artificial Sequence

15

<400> 10

20

gcggatcctt accgagccca agtgcc

<210> 11

25

<211> 27

<212> DNA

<213> Artificial Sequence

30

<400> 11

35

gcggatcctc acttcctttg tggccag

<210> 12

40

<211> 30

<212> DNA

<213> Artificial Sequence

45

<400> 12

50

gttggcgtac caagaccgag cccaagtgcc

<210> 13

55

EP 1 022 332 A1

<211> 1894

<212> DNA

<213> *Scopulariopsis brevicaulis* IF04843

<220>

<221> CDS

<222> (1)... (1721)

<221> intron

<222> (1722)... (1776)

<221> CDS

<222> (1777)... (1894)

<400> 13

caacctacgt ctctgtcaat cgacaattcc acgtatcctt ctatcgacta caactccgcc 60
 cctccaaacc tctcgactct tgccaacaac agcctcttcg agacatggag gccgagggca 120
 cactctcttc cgccccagaa ccagatcggc gatccgigta tgcactacac cgaccccgag 180
 acaggaatct tccacgtcgg ctggctgtac aacggcaatg gcgcttcgg cgccacgacc 240
 gaggatctcg tcacctataa ggatctcaac cccgacggag cgcagatgat ccttcggggt 300
 ggtgtgaalg accccattgc tgtctttgac ggcgcggtta ttcccagtgg catlgaatgg 360
 aaaccaccca tgatgtatac ctcggtgtca tacaigccca tctcctggag catcgcttac 420
 accaggggaa gcgagacca ctctctcgca gtgtcgtccg acggcggtaa gaacttcacc 480
 aagctgggtc agggccccgt catctcttcg cctcccttcg gcgccaacgt gaccagctgg 540
 cgtgacccct tctgtttcca aaacccccag ttcgactctc tcttcgaaag cgagaacggc 600
 acgtgggtaca ccgttatctc tggtaggcac cactgtgacg gcccttcgc gttcctctac 660
 cgtcagcacg accccgactt ccagtactgg gactaccttg gaccgtggtg gaacgaggaa 720
 gggaactcga cctggggcag cggtagctgg gctggccggt ggggctacaa cttcgaggtc 780
 atcaacattg tcggctctga cgatgatggc tacaaccccg acggtgaaat ctttgccacg 840
 gtaggtaccg aatggtcgtt tgaccccatc aaaccgcagg cctcggacaa caggagatg 900
 ctctggggccg cgggcaacat gactctcgag gacggcgata tcaagttcac gccaaagcatg 960
 gcgggctacc tcgactgggg tctatcgccg tatgccgccg ctggcaagga gctgcccgct 1020

EP 1 022 332 A1

tcttcaaagc cttcgagaa gagcgggtgcg ccggaccggt tcgtgtcgta cctgtggctc 1080

5

accggtgact acttcgaggg ccacgacttc cccaccccgcc agcagaattg gaccggctcg 1140

cttttgcttc cgcgtgagct gagcgtcggg acgattccca acgttgtcga caacgagctt 1200

gctcgcgaga cgggctcttg gagggttggc accaaccgaca ctggcgtgct tgagctggtc 1260

10

actctgaagc aggagattgc tcgcgagacg ctggctgaaa tgaccagcgg caactccttc 1320

accgaggcga gcaggaatgt cagctcgccc ggatctaccg ccttccagca gtccctggat 1380

tccaagttct tcgtcctgac cgcctcgctc tccctccctt cgtcggctcg cgactccgac 1440

15

ctcaaggctg gtttcgagat cctgtcgctc gagtttgagt cgaccacggt ctactaccag 1500

ttttccaacg agtccatcat cattgaccgg agcaacitga gtgctgccgc cttgactacc 1560

20

gatggaatcg acacccgcaa cgagtttggc aagaigcgcc tgtttgatgt tgtcgagggt 1620

gaccaggagc glatcgagac gctcgatctc actatigtgg ttgataactc gatcgttgag 1680

gttcatgcca acgggcgatt cgtctcgagc acttgggttc ggtaagtggc acgccgacca 1740

25

cgcttcttaa tttctccaat actaacatta lcacagttcg tggtagagt cgtccaagga 1800

catcaagttc ttccacgatg gcgacagcac ggttcagttc tcgaacatca ccgtctacga 1860

gggactgttt gacgcctggc cggagcgggc cagg 1894

30

<210> 14

<211> 26

35

<212> DNA

<213> Artificial Sequence

40

<400> 14

gcggatccat gaaactctca actgtt

45

<210> 15

<211> 26

50

<212> DNA

<213> Artificial Sequence

55

<400> 15

5 gcggatcctt accgaacca agtgct

<210> 16

10

<211> 25

<212> DNA

15

<213> Artificial Sequence

<400> 16

20

gcggatcctt acctggcccg ctccg

25

<210> 17

<211> 33

<212> DNA

30

<213> Artificial Sequence

35

<400> 17

cgactcgtag cacgaacgaa cccaagtgct cag

40

Claims

45

1. A polypeptide comprising the amino acid sequence of SEQ ID No. 1 or a homologue thereof.

2. A DNA encoding a polypeptide according to Claim 1.

50

3. A DNA according to Claim 2 comprising the nucleotide sequence of SEQ ID No. 2.

4. A polypeptide comprising the amino acid sequence of SEQ ID No. 3 or a homologue thereof.

5. A DNA encoding a polypeptide according to Claim 4.

55

6. A DNA according to Claim 5 comprising the nucleotide sequence of SEQ ID No. 4.

7. A vector comprising a DNA according to Claim 2, 3, 5 or 6.

8. A host cell transformed by a vector according to Claim 7.

9. A process for producing a β -fructofuranosidase comprising the steps of:

5 cultivating a host cell according to Claim 8, and
collecting the β -fructofuranosidase from the host and/or the culture thereof.

10. A process for producing fructooligosaccharides comprising the step of bringing sucrose into contact with a host cell
according to Claim 8 or a β -fructofuranosidase obtained by the process according to Claim 9.

10

15

20

25

30

35

40

45

50

55





FIG. 1A






FIG. 1C

1.8 kbp BamHI DNA FRAGMENT B

1.8 kbp BamHI DNA FRAGMENT A

pY2831




FIG. 1D




FIG. 2 A




FIG. 2B

EP 1 022 332 A1

INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP98/04087

A. CLASSIFICATION OF SUBJECT MATTERInt.Cl⁶ C12N9/10, C12N15/54, C12N1/19, C12P19/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

Int.Cl⁶ C12N9/10, C12N15/54, C12N1/19, C12P19/00

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
DDBJ/EMBL/GenBank**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	L.M. Boddy et al., "Purification and characterization of an <i>Aspergillus niger</i> invertase and its DNA sequence", Curr. Genet., Vol. 24, P.60-66 (1993)	1-10
P, X	WO, 97/34004, A (Meiji Seika Kaisha, Ltd.), 18 September, 1997 (18. 09. 97) (Family: none)	1-10

☐ Further documents are listed in the continuation of Box C.
 ☐ See patent family annex.

* Special categories of cited documents:	T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A" document defining the general state of the art which is not considered to be of particular relevance	X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"E" earlier document but published on or after the international filing date	Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"A" document member of the same patent family
"O" document referring to an oral disclosure, use, exhibition or other means	
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search
27 November, 1998 (27. 11. 98)Date of mailing of the international search report
8 December, 1998 (08. 12. 98)Name and mailing address of the ISA/
Japanese Patent Office

Authorized officer

Facsimile No.

Telephone No.

Form PCT/ISA/210 (second sheet) (July 1992)